

TOP GUN™

SPECIMEN LABEL

Pellet Rodenticide Place Pack with BITREX®

**Mouse and Rat Poison
for Indoor and Outdoor Rodent Control
Kills House Mice, Norway Rats and Roof Rats**

**Kills House Mice, Norway Rats,
and Roof Rats.**

House Mice, Norway Rats and Roof Rats usually consume a lethal dose in a single night's feeding, but it may take two or more days from time of bait consumption for these rodents to die. Rodents cease feeding after consuming a toxic dose. TOP GUN™ Pellet Rodenticide Place Pack is effective against anticoagulant-resistant mice and rats.

ACTIVE INGREDIENTS: Bromethalin:
N-methyl-2,4-dinitro-N-(2,4,6-tribromophenyl)-
6-(trifluoromethyl) benzenamine 0.01%
OTHER INGREDIENTS: 99.99%
TOTAL: 100.00%

EPA REG. NO. 67517-72-56
EPA EST. NO. 56-OH-1

1/2 OZ. PLACE PACK

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Marketed By: J.T. Eaton & Co., Inc.,
1393 East Highland Road, Twinsburg, OH 44087 U.S.A.

TOP GUN™ is a trademark of the
J.T. Eaton & Co., Inc., Twinsburg, OH 44087
BITREX® is a registered trademark of Macfarlan Smith, Scotland

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling bait.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of the throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN:

TOP GUN™ Pellet Rodenticide is not an anticoagulant type of rodenticide. If ingested, limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets. Keep container closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty containers. Dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations

must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.

3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings and similar man-made structures. This product may also be used in alleys located in urban areas, inside transport vehicles (ships, trains, aircraft), and in and around related port or terminal buildings. Do not use TOP GUN™ Pellet Rodenticide Place Pack in sewers. Do not place this bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast place packs or bait.

SELECTION OF TREATMENT AREAS: Determine areas where rodents will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food accessible to rodents as possible.

APPLICATION DIRECTIONS:

The active ingredient in TOP GUN™ Pellet Rodenticide Place Pack is different from anticoagulant rodenticides. Individual mice cease feeding after consuming a toxic dose.

Rats: Apply 2 to 12 place packs per placement location. Space placements at intervals of 15 to 30 feet.

Mice: Apply 1 to 2 place packs per placement location. Space placements at intervals of 8 to 12 feet. Larger placements, up to 6 place packs, may be needed at points of extremely high mouse activity.

Rats and Mice: Adjust the amount of bait applied to the level of rodent feeding expected at each placement location. Highest bait consumption is expected to occur on the first day or two after treatment. After several days, inspect bait placements and replenish bait at sites where there is heavy feeding or where there is continued evidence of rodent activity. Maintain bait for at least one week and continue baiting until all signs of rodent activity have ceased or it becomes clear that the remaining rodents are not attracted to the bait.

Collect and properly dispose of all dead rodents and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Notice of Warranty

J.T. EATON & CO., INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

For Information on this Pesticide Product (Including Health Concerns, Medical Emergencies, or Pesticide Incidents)

Call 1-800-664-9042 or the
National Pesticide Telecommunications Network
at 1-800-658-7378