Prescription Treatment® brand

250 PROPOXUR

Residual Insecticide

KILLS:

Ants, Booklice, Boxelder Bugs, Brown Dog Ticks, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Drugstore Beetles, Earwigs, Elm Leaf Beetles, Flour Beetles, Grain Weevils, Millipedes, Silverfish, Sowbugs, Spiders, Termites and Waterbugs

FOR USE IN AND AROUND:

Apartments, Homes, Hospitals*, Hotels, Food Storage Areas, Meat Packing Plants, Food Processing Plants, Motels, Nursing Homes*, Restaurants and Other Food Handling Establishments, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains and Planes), Warehouses and Other Commercial and Industrial Buildings

[†] Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm.

EPA Reg. No. 499-501

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vorniting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Product contains petroleum distillate - vomiting may cause aspiration pneumonia. Product contains cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow children or pets in treated areas until surfaces are dry. Keep out of reach of children.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks, open flame or pilot lights. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact any person or pet, either directly or indirectly. Keep people and pets out of the area during application. Do not smoke while using. Provide adequate ventilation of area being treated. Do not apply to humans, pets or plants or contaminate feed, foodstuffs, dishes or utensils. Cover and avoid spraying fish aquariums.

HOW TO USE PRESCRIPTION TREATMENT BRAND 250 PROPOXUR

SPOT TREATMENT: Apply to surfaces as a spot treatment only. Do not spray up into air. Point spray opening toward surface to be sprayed and press button firmly. Hold about 12" from the surface being sprayed. Spray around baseboards, doors, window sills and plumbing and other places where insects may enter. Spray into inaccessible areas behind and beneath sinks, stoves, cabinets, refrigerators, damp areas, and in and around waste containers. Avoid excessive wetting of asphalt, tile, rubber and plastic. Do not treat open floor areas, furniture, beds or bedding. Clean any exposed area where food or people may contact.

CRACK & CREVICE® TREATMENT: Use 250 Propoxur with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject



250 Propoxur into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings where insects may enter. Spray into inaccessible areas behind and beneath sinks, stoves, cabinets, refrigerators, damp areas and in and around waste containers.

ENTRY RESTRICTION: Do not allow people or pets to enter the treated area until sprays have dried.

INDOOR TREATMENTS

ANTS: Apply as a spot treatment or Crack & Crevice treatment. Spray into nests, trails, around doors, windows, into structural cracks and crevices, and wherever ants enter house. Spray directly on insects in these locations where possible. Repeat as necessary. Do not treat entire area of floor or floor covering.

CRAWLING PESTS: Cockroaches, Waterbugs, Spiders, Brown Dog Ticks, Earwigs, Crickets, Sowbugs, Millipedes, and Centipedes – Apply as a spot treatment or Crack & Crevice treatment. Spray into hiding places such as cracks and crevices, moist areas, openings around pipes and sinks, under refrigerators, behind baseboards, coffee makers, meter boxes and manholes. Spray directly on insects in these locations where possible. Repeat as necessary.

BOOKLICE & SILVERFISH: Spray cracks and crevices in baseboards, doorframes, bookcases and storage areas. Spray directly on insects in these locations where possible. Repeat as necessary.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUGSTORE BEETLES (in exposed adult and larval stages) and hibernating stages of BOXELDER BUGS, ELM LEAF BEETLES, CLUSTER FLIES and CLOVER MITES: Contact as many insects as possible when treating infested areas such as cracks and crevices in wall voids, around windows and doorframes, under siding, in machinery, and in dark, secluded areas. Repeat as necessary.

Localized treatment of wood destroying insects: CAR-PENTER ANTS and CARPENTER BEES — Locate source of infestation. Inject 5 – 10 seconds of spray into insect tunnels and cavities. For CARPENTER BEES, seal entrance openings with small pieces of stainless steel wool after spray application. FOR TERMITES: Locate source of infestation. Apply to voids and channels in damaged wooden members of a structure or to cracks, spaces or bearing joints between wooden members of a structure. Inject 5 - 10 seconds spray into insect tunnels and cavities. This application is not a substitute for mechanical alteration, soil treatment or foundation treatment but is merely a supplemental treatment. This product is intended to kill workers or winged reproductive forms which may be present in the treated areas.

APPLICATIONS IN AIRCRAFT: This product can be applied in cargo areas but not in aircraft cabins.

APPLICATION IN FOOD HANDLING ESTABLISHMENTS

Food handling establishments are places other than private residences in which food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK & CREVICE TREATMENT ONLY: Includes areas for receiving, storage, packaging, (canning bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) are also considered a food area.

Apply prescribed amounts of 250 Propoxur as a Crack & Crevice treatment using the supplied injection tube or other Whitmire Micro-Gen equipment in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces.

APPLICATION OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK & CREVICE TREATMENT IS NOT PERMITTED.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store away from heat or open flames, in an area inaccessible to children.

PESTICIDE DISPOSAL: Waste resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product in accordance with the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly full, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com
© 2006 Whitmire Micro-Gen Research Laboratories, Inc.